

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 460 332 A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 90440051.2

(51) Int. Cl.⁵: A01N 3/00

(22) Date of filing: 07.06.90

(43) Date of publication of application:
11.12.91 Bulletin 91/50

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(71) Applicant: Sellegaard, Lars E.
9 Domaine du Camp Lauvas
F-06250 Mougins(FR)

(72) Inventor: Sellegaard, Lars E.
9 Domaine du Camp Lauvas
F-06250 Mougins(FR)

(74) Representative: Bossard, Jacques-René
Cabinet MEYER & COURTASSOL Bureau
EUROPE 20 Place des Halles
F-67000 Strasbourg(FR)

(54) Method for preserving flowers particularly roses.

(57) Method for preserving natural flowers, in which the lower end of the stem of the flower is immersed into a water/glycerol solution of a dry composition comprising mineral salts, citric acid and dyes, characterized in that the flower to be treated is severed from the upper end of its stem so that the remaining section of said stem still connected to the flower is shorter than 5 cm and preferably of about 1 cm so that the path of said solution towards the flower be as short as possible.

EP 0 460 332 A1

In prior patents (F-85-11992 and F-85 16264) the applicant has described a process and a composition for treating plants in view of keeping their aspect and their natural freshness during an extended period of time after being extracted from their natural environment. The process consisted in immersing the newly cut lower portion of the stem of the plant into a specific solution containing a number of mineral salts and of dyes in a mixture of water and glycerol during a given duration and at a given temperature.

This process and this composition were effective in keeping the freshness of many plants and bushes : after treatment the stems and leaves of said plants kept during many months and even years the aspect and the touch of the fresh plants before treatment.

One critical feature of said compositions was its content in citric acid, which was of 4-5 %, allowing a proportion of glycerol up to 40 % and a treatment temperature of 38-45° C, said conditions being more favourable than the conditions of still older known similar processes. However, even with said conditions, this process was still difficult to work in case of some flowers such as roses, having specially delicate petals as well as a fragile structure such as the overall shape and even a transient scent.

The further works of the Applicant have reached to a modification of this process specially adapted to this specific product and more specially to the treatment of roses.

The invention has therefore as a first object a process of the above general nature but adapted to the preservation treatment of fragile flowers and more specifically of roses. It has also for object a dry composition specially adapted to the use in this process and a liquid solution obtained from said composition. It has also for object the flowers the freshness, the aspect and the scent thereof have been preserved by the use of said process, and more specially the roses.

A critical feature of this process is that the path of the solution up to the flower to be treated should be as short as possible, i.e shorter than 5 cm and preferably of less. To this effect, the flower is severed as high as possible at the upper end of its stem and the small remaining section is immersed into the solution avoiding any contact of the petals with said solution. At the end of the overall treatment as it will be described hereunder, the stem with its leaves, separately treated by the usual process, is connected to the flower through a glued rod or pin, e.g of plastic.

Another feature of this invention is the dry composition from which the treatment solution is prepared. In fact, the constituents of this composition are similar to those of the prior known compositions, as described in the two French patents mentioned above, but the proportions thereof have been adapted to the specific treatment of roses.

The following table shows said compositions :

COMPONENT	PROPORTION RANGES %	PREFERRED PROPORTION %
Potassium Nitrate	25-40	30
Sodium Sulphate	2-6	4
Manganèse Sulphate monohyd.	6-10	8
Calcium hydrogen Phosphate	2-6	4
Citric Acid	2-5	4

At least one of the following dyes making the remaining 30-60 %, preferably 50 % :

Tartrazine (E 102)
 Patent Blue (E131)
 Patent Green (ES 11 or FD & C N° 3)
 Amaranth Red (E 123)
 Quinoline Yellow (E 104)
 Sunset Yellow (E 110)
 Ponceau 4 R (E 124)

Wool Green (E 142)

The treatment solution comprises about 25g/l of said dry composition into a solvent comprising a mixture water/glycerol in a volume proportion between 75/25 and 55/45.

The process of the invention in its application to roses comprises the following steps :

- 5 1. Sever the rose at the upper end of its stem, the remaining section being shorter than 5 cm and preferably of about 1 cm.
2. Immerse said section into the solution, said solution being slowly circulated in a perforated tube serving to support the flower, the temperature of said solution being between 33 and 44 ° C.
3. Continue said immersion during 3-7 days at an ambient temperature of 22-28 ° C and at an ambient
- 10 humidity less than 65 %
4. At the end of this period, hang the treated flower upside down during about 4 weeks at an ambient temperature between 20 and 30 ° C and a relative humidity less than 60 %.
5. Connect the treated flower with a stem, preferably a treated stem with leaves, through a glued rod or pin.
- 15 The treated roses fully keep their freshness and their scent during a very long period, of at least one year.

To obtain special visual effects, the stem which is assembled to the treated flower can be itself treated with the same process but using a different dye so as the leaves are of a different color.

It is clear that this process is specially directed to the treatment of roses but can be used for any

20 flowers having preferably strong petals such as those forming the families of MATRICARIA and PROTEA and also different wild flowers of the APIACEAE family.

Claims

- 25 1. Method for preserving natural flowers, in which the lower end of the stem of the flower is immersed into a water/glycerol solution of a dry composition comprising mineral salts, citric acid and dyes, characterized in that the flower to be treated is severed from the upper end of its stem so that the remaining section of said stem still connected to the flower is shorter than 5 cm and preferably of about 1 cm so that the path of said solution towards the flower be as short as possible.
- 30 2. Method according to claim 1, in which said dry composition comprises 25-40 %, preferably 30 % of potassium nitrate, 2-6 %, preferably 4 % of sodium sulphate, 6-10 %, preferably 8 % of manganese sulphate monohydrate, 2-6 %, preferably 4 % of calcium hydrogen phosphate and 2-5 %, preferably 4 % of citric acid, and the remaining 30-60 %, preferably 50 % comprising at least one dye selected
- 35 amongst :

Tartrazine	(E 102)
Patent Blue	(E131)
Patent Green	(ES 11 or FD & C N ° 3)
Amaranth Red	(E 123)
40 Quinoline Yellow	(E 104)
Sunset Yellow	(E 110)
Ponceau 4 R	(E 124)
Wool Green	(E 142)
- 45 3. Method according to claim 2, in which the treatment solution comprises about 25g/l of said dry composition dissolved in a mixture water/glycerol in a proportion between 75/25 and 55/45.
4. Method according to any of claims 1 to 3, in which said remaining section of the stem is immersed into said solution at a temperature between 33 and 44 ° C during 3 to 7 days, at an ambient temperature
- 50 between 22 and 28 ° C and at an ambient humidity less than 65 %.
5. Method according to any of the preceding claims, in which, at the end of said treatment by said solution, the flower is kept upside down during about 4 weeks, at a temperature comprised between 20 and 30 ° C, and at a relative humidity less than 60 %, then assembled with a stem also previously
- 55 treated by the same method.
6. Method according to claim 5, in which said assembly is made by glueing a pin inserted at its both ends into the remaining stem of the flower and into the stem to be assembled.

7. Method according to claims 5 and 6, in which the stem to be assembled has been treated by a solution containing a different dye so as to obtain leaves of a different color.
8. Method according to any of the preceding claims, in which the solution is contained in a perforated tube receiving the stem section and in which said solution constantly slowly circulated.
9. Method according to any of the preceding claims, in which the flower to be treated is a rose.
10. Dry composition for use in the method of any of claims 1 to 9, comprising 25-40 %, preferably 30 % of potassium nitrate, 2-6, preferably 4 % of sodium sulphate, 6-10 %, preferably 8 % of manganese sulphate monohydrate, 2-6 %, preferably 4 % of calcium hydrogen phosphate and 2-5 %, preferably 4 % of citric acid, and the remaining 30-60 %, preferably 50 % comprising at least one dye selected amongst :

Tartrazine	(E 102)
Patent Blue	(E131)
Patent Green	(ES 11 or FD & C N° 3)
Amaranth Red	(E 123)
Quinoline Yellow	(E 104)
Sunset Yellow	(E 110)
Ponceau 4 R	(E 124) Wool Green (E 142)
11. Solution for use in the method of any of claims 1 to 9, comprising about 25 g of the dry composition of claim 10 dissolved in one liter of a mixture water/glycerol in a proportion comprised between 75/25 and 55/45
12. Flowers and specially roses treated by the method of any of the preceding claims, for preserving its freshness, its aspect and its scent during a long period of time, at least one year.



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 90 44 0051

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
D,X	GB-A-2 178 294 (L.E. SELLEGAARD)	10,11	A 01 N 3/00
D,A	* Page 2, lines 37-62; claims *	1-9,12	
A	US-A-4 349 459 (C. ROMERO-SIERRA et al.)		
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			A 01 N
The present search report has been drawn up for all claims			
Place of search		Date of completion of search	Examiner
The Hague		11 February 91	DONOVAN T.M.
CATEGORY OF CITED DOCUMENTS			
X: particularly relevant if taken alone			
Y: particularly relevant if combined with another document of the same category			
A: technological background			
O: non-written disclosure			
P: intermediate document			
T: theory or principle underlying the invention			
E: earlier patent document, but published on, or after the filing date			
D: document cited in the application			
L: document cited for other reasons			
&: member of the same patent family, corresponding document			

THIS PAGE BLANK (USPTO)